

09/863,976

=> d 12 abs ibib kwic 26 29

L2 ANSWER 26 OF 33 HCAPLUS COPYRIGHT 2004 ACS on STN
AB A method for **stabilizing** intracellular proteins known to cause disease, said method comprising contacting a cell with **stabilizing** agents such as DMSO, sugars, amino acids and TMAO (trimethylamine N-oxide), wherein the proteins are **stabilized** and the disease state lessened.
ACCESSION NUMBER: 1997:684267 HCAPLUS
DOCUMENT NUMBER: 127:341811
TITLE: Correction of genetic defects using chemical **chaperones**
INVENTOR(S): Welch, William J.; Brown, C. Randell; Tatzelt, Jorg
PATENT ASSIGNEE(S): Regents of the University of California, USA; Welch, William J.; Brown, C. Randell; Tatzelt, Jorg
SOURCE: PCT Int. Appl., 86 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9737645	A1	19971016	WO 1997-US5846	19970409
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2250780	AA	19971016	CA 1997-2250780	19970409
AU 9724485	A1	19971029	AU 1997-24485	19970409
AU 734905	B2	20010628		
EP 959877	A1	19991201	EP 1997-920243	19970409
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2000509971	T2	20000808	JP 1997-536453	19970409
US 2001021500	A1	20010913	US 2001-823657	20010330
US 6541195	B2	20030401		

PRIORITY APPLN. INFO.:

US 1996-15155P P 19960410
US 1997-838691 A2 19970409
WO 1997-US5846 W 19970409
US 1999-291406 A1 19990413

TI Correction of genetic defects using chemical **chaperones**
AB A method for **stabilizing** intracellular proteins known to cause disease, said method comprising contacting a cell with **stabilizing** agents such as DMSO, sugars, amino acids and TMAO (trimethylamine N-oxide), wherein the proteins are **stabilized** and the disease state lessened.
ST genetic disease **conformation** therapy **chaperone** chem
IT Brain, disease
Prion diseases
(Creutzfeldt-Jakob; correction of genetic defects using chemical **chaperones**)
IT Lipoprotein receptors

DELACROIX

- RL: ADV (Adverse effect, including toxicity); BPR (Biological process);
BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological
study); PROC (Process)
(LDL, **mutant**; correction of genetic defects using chemical
chaperones)
- IT Prion proteins
RL: ADV (Adverse effect, including toxicity); BPR (Biological process);
BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological
study); PROC (Process)
(PrPc, **mutant**; correction of genetic defects using chemical
chaperones)
- IT Prion proteins
RL: ADV (Adverse effect, including toxicity); BPR (Biological process);
BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological
study); PROC (Process)
(PrPSc, **mutant**; correction of genetic defects using chemical
chaperones)
- IT Gangliosidosis
(Tay-Sachs disease; correction of genetic defects using chemical
chaperones)
- IT Nervous system
(amyotrophic lateral sclerosis; correction of genetic defects using
chemical **chaperones**)
- IT Liver, disease
(chronic; correction of genetic defects using chemical **chaperones**
)
- IT Alzheimer's disease
Antitumor agents
Cataract
Emphysema
Marfan syndrome
Neoplasm
Scurvy
Stabilizing agents
(correction of genetic defects using chemical **chaperones**)
- IT **Chaperonins**
RL: BAC (Biological activity or effector, except adverse); BPR (Biological
process); BSU (Biological study, unclassified); THU (Therapeutic use);
BIOL (Biological study); PROC (Process); USES (Uses)
(correction of genetic defects using chemical **chaperones**)
- IT Amino acids, biological studies
Carbohydrates, biological studies
Polyoxyalkylenes, biological studies
RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
(Uses)
(correction of genetic defects using chemical **chaperones**)
- IT Nervous system
(disease, Gerstmann-Straussler syndrome; correction of genetic defects
using chemical **chaperones**)
- IT Hypercholesterolemia
(familial; correction of genetic defects using chemical **chaperones**
)
- IT Brain, disease
Prion diseases
(fatal familial insomnia; correction of genetic defects using chemical
chaperones)
- IT Insomnia
(fatal familial; correction of genetic defects using chemical

- chaperones)**
- IT Disease, animal
(genetic, protein **conformation** defects from; correction of genetic defects using chemical **chaperones**)
- IT Endocrine system
(leprechaunism; correction of genetic defects using chemical **chaperones**)
- IT Mental disorder
(maple syrup urine disease; correction of genetic defects using chemical **chaperones**)
- IT CFTR (cystic fibrosis transmembrane conductance regulator)
Collagens, biological studies
Crystallins
Fibrillins
Insulin receptors
Prion proteins
Rhodopsins
p53 (protein)
RL: ADV (Adverse effect, including toxicity); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(**mutant**; correction of genetic defects using chemical **chaperones**)
- IT Bone, disease
(osteogenesis imperfecta; correction of genetic defects using chemical **chaperones**)
- IT Alcohols, biological studies
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(polyhydric; correction of genetic defects using chemical **chaperones**)
- IT Collagens, biological studies
RL: ADV (Adverse effect, including toxicity); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(procollagens, type I, pro- α , **mutant**; correction of genetic defects using chemical **chaperones**)
- IT **Conformation**
(protein; correction of genetic defects using chemical **chaperones**)
- IT Eye, disease
(retinitis pigmentosa; correction of genetic defects using chemical **chaperones**)
- IT Brain, disease
Prion diseases
(scrapie; correction of genetic defects using chemical **chaperones**)
- IT Brain, disease
(spongiform encephalopathy; correction of genetic defects using chemical **chaperones**)
- IT Amyloid
RL: ADV (Adverse effect, including toxicity); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(β -; correction of genetic defects using chemical **chaperones**)
- IT Animal cell line
(AF508; correction of genetic defects using chemical

chaperones)

- IT 9067-96-3, α -Ketoacid dehydrogenase
 RL: ADV (Adverse effect, including toxicity); BPR (Biological process);
 BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological
 study); PROC (Process)
 (correction of genetic defects using chemical **chaperones**)
- IT 56-12-2, Gaba, biological studies 56-40-6, Glycine, biological studies
 56-41-7, Alanine, biological studies 56-81-5, Glycerol, biological
 studies 56-86-0, Glutamic acid, biological studies 67-68-5, Dmsol,
 biological studies 87-89-8, Inositol 99-20-7D, Trehalose,
 isofluorolol 107-35-7, Taurine 107-43-7, Betaine 107-97-1,
 Sarcosine 147-85-3, Proline, biological studies 149-32-6, Erythritol
 1184-78-7, Trimethylamine N-oxide 7789-20-0, Water-d2 25322-68-3,
 Polyethylene glycol 34522-32-2, Octopine
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological
 study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES
 (Uses)
 (correction of genetic defects using chemical **chaperones**)
- IT 302-95-4, Sodium deoxycholate 9002-93-1, Triton x-100
 RL: NUU (Other use, unclassified); USES (Uses)
 (correction of genetic defects using chemical **chaperones**)
- IT 9012-33-3 9041-92-3, α 1 Antitrypsin 9054-89-1, Superoxide
 dismutase
 RL: ADV (Adverse effect, including toxicity); BPR (Biological process);
 BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological
 study); PROC (Process)
 (**mutant**; correction of genetic defects using chemical
chaperones)

L2 ANSWER 29 OF 33 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN

AB Recently, we found that different low molecular weight compounds, all
 known to **stabilize** proteins in their native **conformation**
 , are effective in correcting the temperature-sensitive protein folding
 defect associated with the Δ F508 cystic fibrosis transmembrane
 regulator (CFTR) protein. Here we examined whether the folding of other
 proteins which exhibit temperature- sensitive folding defects also could
 be corrected via a similar strategy. Cell lines expressing
 temperature-sensitive **mutants** of the tumor suppressor protein
p53, the viral oncogene protein pp60(src), or a ubiquitin
 activating enzyme E1, were incubated at the nonpermissive temperature
 (39.5°C) in the presence of glycerol, trimethylamine N-oxide or
 deuterated water. In each case, the cells exhibited phenotypes similar to
 those observed when the cells were incubated at the permissive temperature
 (32.5°C), indicative that the particular protein folding defect had
 been corrected. These observations, coupled with our earlier work and much
 older studies in yeast and bacteria, indicate that protein
stabilizing agents are effective in vivo for correcting protein
 folding abnormalities. We suggest that this type of approach may prove to
 be useful for correcting certain protein folding abnormalities associated
 with human diseases.

ACCESSION NUMBER: 97099314 EMBASE
 DOCUMENT NUMBER: 1997099314
 TITLE: Correcting temperature-sensitive protein folding defects.
 AUTHOR: Brown C.R.; Hong-Brown L.Q.; Welch W.J.
 CORPORATE SOURCE: C.R. Brown, UCSF, UC Box 0854, San Francisco, CA 94143,
 United States. crb@itsa.ucsf.edu
 SOURCE: Journal of Clinical Investigation, (1997) 99/6 (1432-1444).
 Refs: 36

09/863,976

ISSN: 0021-9738 CODEN: JCINAO

COUNTRY: United States

DOCUMENT TYPE: Journal; Article

FILE SEGMENT: 022 Human Genetics

029 Clinical Biochemistry

LANGUAGE: English

SUMMARY LANGUAGE: English

AB Recently, we found that different low molecular weight compounds, all known to **stabilize** proteins in their native **conformation**, are effective in correcting the temperature-sensitive protein folding defect associated with the $\Delta F508$ cystic fibrosis transmembrane regulator (CFTR) protein. Here. . . other proteins which exhibit temperature-sensitive folding defects also could be corrected via a similar strategy. Cell lines expressing temperature-sensitive **mutants** of the tumor suppressor protein **p53**, the viral oncogene protein pp60(src), or a ubiquitin activating enzyme E1, were incubated at the nonpermissive temperature (39.5°C) in the. . . been corrected. These observations, coupled with our earlier work and much older studies in yeast and bacteria, indicate that protein **stabilizing** agents are effective in vivo for correcting protein folding abnormalities. We suggest that this type of approach may prove to.

CT Medical Descriptors:

*protein folding

*temperature sensitive mutant

article

cell growth

cell line

cell structure

gene expression

immunoblotting

priority journal

chaperone: EC, endogenous compound

protein kinase p60: EC, endogenous compound

protein p53: EC, endogenous compound

ubiquitin protein ligase: EC, endogenous compound

=>

09/863,976

FILE 'HCAPLUS' ENTERED AT 22:06:56 ON 24 MAY 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 22:06:56 ON 24 MAY 2004
COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 22:06:56 ON 24 MAY 2004
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

FILE 'MEDLINE' ENTERED AT 22:06:56 ON 24 MAY 2004

=> s p53 and (conformation? or stabili?) and chaperon? and (mutant? or
wild(2a)type?)

L1 85 P53 AND (CONFORMATION? OR STABILI?) AND CHAPERON? AND (MUTANT?
OR WILD(2A) TYPE?)

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 33 DUP REM L1 (52 DUPLICATES REMOVED)

=> d l2 ibib 1-33

09/863,976

(FILE 'HOME' ENTERED AT 22:06:34 ON 24 MAY 2004)

FILE 'HCAPLUS, BIOSIS, EMBASE, MEDLINE' ENTERED AT 22:06:56 ON 24 MAY 2004

L1 85 S P53 AND (CONFORMATION? OR STABILIZ?) AND CHAPERON? AND (MUTANT
L2 33 DUP REM L1 (52 DUPLICATES REMOVED)

=> s l1 and spectroscop?

L3 3 L1 AND SPECTROSCOP?

=> d l3 abs ibib kwic 1-3

L3 ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2004 ACS on STN

AB We disclose a method of **stabilizing** the native state of a polypeptide, the method comprising exposing the polypeptide to a **stabilizing** mol. capable of binding to the polypeptide at a site which at least partially overlaps a functional site in its native state. The authors describe the isolation and identification of a **stabilizing** peptide CDB3, which is capable of binding the tumor suppressor protein **p53** near its DNA binding site, and **stabilizing** the native form of the protein. Since the binding of DNA itself **stabilizes** **p53** core domain, and it binds very tightly, **stabilization** by a peptide such as CDB3 is needed only for **mutants** where DNA binding is impaired because **mutant p53** is in denatured **conformation**. Once the protein has bound DNA, the peptide is not needed any more. The ability of CDB3 to induce refolding of **p53** core domain, together with the observation that DNA can displace it from **p53**, led the authors to propose the a "**chaperone**" mechanism for rescuing a denatured oncogenic protein: CDB3 binds only the native state of the oncogenic protein which is able to bind DNA, probably immediately on biosynthesis, and therefore shifts the equilibrium towards the native state. Then DNA can bind the protein, displacing the peptide, which is free again to bind another protein mol. Exemplary design of potential **P53** core domain binding peptides, screening of the CDB peptides for binding **p53** core domain, identification of the **P53** core domain binding peptide CDB3, characterization of CDB3-**P53** core domain binding and binding of fluorescein-labeled CDB3 are reported. **Stabilizing** mols. and/or compns. of the invention can be used in the treatment of any animal or human disease where errors in protein **conformation**, folding and aggregation contribute to the disease. Examples include cancer, cystic fibrosis and neuro-degeneration. In a particularly preferred embodiment, the disease is cancer.

ACCESSION NUMBER: 2003:133296 HCAPLUS

DOCUMENT NUMBER: 138:166255

TITLE: **Stabilization** of the native **conformation** of a **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomolecules and application to treatment of cancer and other diseases

INVENTOR(S): Friedler, Assaf; Fersht, Alan

PATENT ASSIGNEE(S): Medical Research Council, UK

SOURCE: PCT Int. Appl., 73 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

DELACROIX

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003014144	A2	20030220	WO 2002-GB3668	20020809
WO 2003014144	A3	20031127		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1414846	A2	20040506	EP 2002-749128	20020809
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
PRIORITY APPLN. INFO.:				
			GB 2001-19557	A 20010810
			GB 2001-27917	A 20011121
			GB 2002-10740	A 20020510
			WO 2002-GB3668	W 20020809

TI **Stabilization** of the native **conformation** of a **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomolecules and application to treatment of cancer and other diseases

AB We disclose a method of **stabilizing** the native state of a polypeptide, the method comprising exposing the polypeptide to a **stabilizing** mol. capable of binding to the polypeptide at a site which at least partially overlaps a functional site in its native state. The authors describe the isolation and identification of a **stabilizing** peptide CDB3, which is capable of binding the tumor suppressor protein **p53** near its DNA binding site, and **stabilizing** the native form of the protein. Since the binding of DNA itself **stabilizes** **p53** core domain, and it binds very tightly, **stabilization** by a peptide such as CDB3 is needed only for **mutants** where DNA binding is impaired because **mutant p53** is in denatured **conformation**. Once the protein has bound DNA, the peptide is not needed any more. The ability of CDB3 to induce refolding of **p53** core domain, together with the observation that DNA can displace it from **p53**, led the authors to propose the a "**chaperone**" mechanism for rescuing a denatured oncogenic protein: CDB3 binds only the native state of the oncogenic protein which is able to bind DNA, probably immediately on biosynthesis, and therefore shifts the equilibrium towards the native state. Then DNA can bind the protein, displacing the peptide, which is free again to bind another protein mol. Exemplary design of potential **P53** core domain binding peptides, screening of the CDB peptides for binding **p53** core domain, identification of the **P53** core domain binding peptide CDB3, characterization of CDB3-**P53** core domain binding and binding of fluorescein-labeled CDB3 are reported. **Stabilizing** mols. and/or compns. of the invention can be used in the treatment of any animal or human disease where errors in protein **conformation**, folding and aggregation contribute to the disease. Examples include cancer, cystic fibrosis and neuro-degeneration. In a particularly preferred embodiment, the disease is cancer.

ST tumor suppressor protein **p53** mutation **conformation** CDB3 peptide anticancer; protein **conformation** mutation

- stabilization** biomol human disease treatment
- IT Enzyme functional sites
 - (active, **stabilizing** biomol. binding to;
 - stabilization** of native **conformation** of human
 - mutant** tumor suppressor protein **p53** and other
 - mutant** proteins using CDB3 peptide and other biomols. and
 - application to treatment of cancer and other diseases)
- IT Fluorescence
 - (anisotropy, biomol. binding detection using; **stabilization**
 - of native **conformation** of human **mutant** tumor
 - suppressor protein **p53** and other **mutant** proteins
 - using CDB3 peptide and other biomols. and application to treatment of
 - diseases)
- IT Differential scanning calorimetry
 - NMR **spectroscopy**
 - Surface plasmon resonance
 - (biomol. binding detection using; **stabilization** of native
 - conformation** of human **mutant** tumor suppressor protein
 - p53** and other **mutant** proteins using CDB3 peptide and
 - other biomols. and application to treatment of cancer and other
 - diseases)
- IT Chromophores
 - Fluorescent substances
 - (biomol. derivatized with; **stabilization** of native
 - conformation** of human **mutant** tumor suppressor protein
 - p53** and other **mutant** proteins using CDB3 peptide and
 - other biomols. and application to treatment of cancer and other
 - diseases)
- IT Amides, biological studies
 - Amines, biological studies
 - Phosphates, biological studies
 - Sulfates, biological studies
 - Sulfides, biological studies
 - RL: BUU (Biological use, unclassified); THU (Therapeutic use); BIOL
 - (Biological study); USES (Uses)
 - (biomol. derivatized with; **stabilization** of native
 - conformation** of human **mutant** tumor suppressor protein
 - p53** and other **mutant** proteins using CDB3 peptide and
 - other biomols. and application to treatment of cancer and other
 - diseases)
- IT Molecular association
 - (biomol.-protein native state; **stabilization** of native
 - conformation** of human **mutant** tumor suppressor protein
 - p53** and other **mutant** proteins using CDB3 peptide and
 - other biomols. and application to treatment of cancer and other
 - diseases)
- IT NMR **spectroscopy**
 - (heteronuclear double resonance, biomol. binding detection using;
 - stabilization** of native **conformation** of human
 - mutant** tumor suppressor protein **p53** and other
 - mutant** proteins using CDB3 peptide and other biomols. and
 - application to treatment of cancer and other diseases)
- IT Protein folding
 - (induction of refolding; **stabilization** of native
 - conformation** of human **mutant** tumor suppressor protein
 - p53** and other **mutant** proteins using CDB3 peptide and
 - other biomols. and application to treatment of cancer and other
 - diseases)
- IT Apoptosis

- (induction of; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Carbohydrates, biological studies
 DNA
 Glycoproteins
 Nucleic acids
 Oligonucleotides
 Peptide nucleic acids
 Peptides, biological studies
 RNA
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (mutant proteins **stabilization** by;
stabilization of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Proteins
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 --(oncogenic, **stabilization** of; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Dissociation constant
 (p53-CDB3; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Disease, animal
 (protein **conformation** error-related; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT **Conformation**
 (protein, **stabilization** of; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Denaturation
 (protein; **stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other **mutant** proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Antitumor agents
 Biochemical molecules
 Drug screening
 Human
 Mutation
 Neoplasm
 Protein engineering
 (**stabilization** of native **conformation** of human **mutant** tumor suppressor protein **p53** and other

- mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT **p53** (protein)
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Proteins
 RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (stabilization of; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Proteins
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (tumor suppressor, stabilization of; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT Phenotypes
 (wild-type, restoring of; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT 497259-83-3P
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (CDB3; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT 56-45-1, L-Serine, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (G245S and R249S **p53** mutations; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT 72-19-5, L-Threonine, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (I195T **p53** mutation; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)
- IT 71-00-1, L-Histidine, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (R175H and R273H **p53** mutations; stabilization of native conformation of human mutant tumor suppressor protein **p53** and other mutant proteins using CDB3 peptide and other biomols. and application to treatment of cancer and other diseases)

- cancer and other diseases)
- IT 56-85-9, L-Glutamine, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (R248Q **p53** mutation; **stabilization** of native
conformation of human **mutant** tumor suppressor protein
p53 and other **mutant** proteins using CDB3 peptide and
 other biomols. and application to treatment of cancer and other
 diseases)
- IT 73-22-3, L-Tryptophan, biological studies
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (R282W **p53** mutation; **stabilization** of native
conformation of human **mutant** tumor suppressor protein
p53 and other **mutant** proteins using CDB3 peptide and
 other biomols. and application to treatment of cancer and other
 diseases)
- IT 58-85-5, Biotin 2321-07-5, Fluorescein
 RL: BUU (Biological use, unclassified); THU (Therapeutic use); BIOL
 (Biological study); USES (Uses)
 (biomol. derivatized with; **stabilization** of native
conformation of human **mutant** tumor suppressor protein
p53 and other **mutant** proteins using CDB3 peptide and
 other biomols. and application to treatment of cancer and other
 diseases)
- IT 497259-83-3DP, fluorescein labeled
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified);
 THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
 (Uses)
 (**stabilization** of native **conformation** of human
mutant tumor suppressor protein **p53** and other
mutant proteins using CDB3 peptide and other biomols. and
 application to treatment of cancer and other diseases)
- L3 ANSWER 2 OF 3 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
- AB **Conformationally** compromised oncogenic **mutants** of the
 tumor suppressor protein **p53** can, in principle, be rescued by
 small molecules that bind the native, but not the denatured state. We
 describe a strategy for the rational search for such molecules. A
 nine-residue peptide, CDB3, which was derived from a **p53** binding
 protein, binds to **p53** core domain and **stabilizes** it in
 vitro. NMR studies showed that CDB3 bound to **p53** at the edge of
 the DNA binding site, partly overlapping it. The fluorescein-labeled
 peptide, FL-CDB3, binds **wild-type p53** core
 domain with a dissociation constant of 0.5 μ M, and raises the apparent
 melting temperatures of **wild-type** and a representative
 oncogenic **mutant**, R249S core domain, gadd45 DNA competes with
 CDB3 and displaces it from its binding site. But this competition does not
 preclude CDB3 from being a lead compound, CDB3 may act as a "
chaperone" that maintains existing or newly synthesized
 destabilized **p53 mutants** in a native
conformation and then allows transfer to specific DNA, which binds
 more tightly. Indeed, CDB3 restored specific DNA binding activity to a
 highly destabilized **mutant** 1195T to close to that of
wild-type level.
- ACCESSION NUMBER: 2002045792 EMBASE
 TITLE: A peptide that binds and **stabilizes p53**
 core domain: **Chaperone** strategy for rescue of
 oncogenic **mutants**.
 AUTHOR: Friedler A.; Hansson L.O.; Veprintsev D.B.; Freund S.M.V.;

09/863,976

Ripplin T.M.; Nikolova P.V.; Proctor M.R.; Rudiger S.;
Fersht A.R.

CORPORATE SOURCE: A.R. Fersht, Cambridge Univ. Chemical Laboratory, Cambridge
Center for Protein Eng., Medical Research Council Centre,
Hills Road, Cambridge CB2 2QH, United Kingdom.
arf25@cam.ac.uk

SOURCE: Proceedings of the National Academy of Sciences of the
United States of America, (22 Jan 2002) 99/2 (937-942).
Refs: 23
ISSN: 0027-8424 CODEN: PNASA6

COUNTRY: United States

DOCUMENT TYPE: Journal; Article

FILE SEGMENT: 016 Cancer
030 Pharmacology
037 Drug Literature Index

LANGUAGE: English

SUMMARY LANGUAGE: English

TI A peptide that binds and **stabilizes p53** core domain:
Chaperone strategy for rescue of oncogenic **mutants**.

AB **Conformationally** compromised oncogenic **mutants** of the
tumor suppressor protein **p53** can, in principle, be rescued by
small molecules that bind the native, but not the denatured state. We
describe a strategy for the rational search for such molecules. A
nine-residue peptide, CDB3, which was derived from a **p53** binding
protein, binds to **p53** core domain and **stabilizes** it in
vitro. NMR studies showed that CDB3 bound to **p53** at the edge of
the DNA binding site, partly overlapping it. The fluorescein-labeled
peptide, FL-CDB3, binds **wild-type p53** core
domain with a dissociation constant of 0.5 μ M, and raises the apparent
melting temperatures of **wild-type** and a representative
oncogenic **mutant**, R249S core domain, gadd45 DNA competes with
CDB3 and displaces it from its binding site. But this competition does not
preclude CDB3 from being a lead compound, CDB3 may act as a "
chaperone" that maintains existing or newly synthesized
destabilized **p53 mutants** in a native
conformation and then allows transfer to specific DNA, which binds
more tightly. Indeed, CDB3 restored specific DNA binding activity to a
highly destabilized **mutant** l195T to close to that of
wild-type level.

CT Medical Descriptors:
*oncogene
*protein DNA binding
*antineoplastic activity
protein domain
in vitro study
nuclear magnetic resonance spectroscopy
dissociation constant
melting point
binding site
protein conformation
surface plasmon resonance
fluorescence
anisotropy
differential scanning calorimetry
drug mechanism
human
article
priority journal
***protein p53**

DELACROIX

09/863,976

*chaperone: DV, drug development
*chaperone: PD, pharmacology
*protein cdb3: DV, drug development
*protein cdb3: PD, pharmacology
*DNA binding protein
DNA: EC, endogenous compound
fluorescein
synthetic peptide: DV, drug development
synthetic peptide: PD, . . .

L3 ANSWER 3 OF 3 MEDLINE on STN

AB The molecular **chaperone** Hsp90 sequesters oncogenic **mutants** of the tumor suppressor **p53** that have unstable core domains. It is not known whether **p53** is bound in an unfolded, partly folded, or distorted structure, as is unknown for the structure of any bound substrate of Hsp90. It is a particularly difficult problem to analyze in detail the structures of large complexes in which one component is (partly) unfolded. We have shown by transverse relaxation-optimized NMR **spectroscopy** combined with cross-correlated relaxation-enhanced polarization transfer (CRINEPT-TROSY) that **p53** core domain bound in an approximately 200-kDa complex with Hsp90 was predominantly unfolded lacking helical or sheet secondary structure. This mode of binding might be a general feature of substrates of Hsp90.

ACCESSION NUMBER: 2002432245 MEDLINE

DOCUMENT NUMBER: PubMed ID: 12163643

TITLE: CRINEPT-TROSY NMR reveals **p53** core domain bound in an unfolded form to the **chaperone** Hsp90.

AUTHOR: Rudiger Stefan; Freund Stefan M V; Veprintsev Dmitry B; Fersht Alan R

CORPORATE SOURCE: Cambridge Centre for Protein Engineering, Cambridge University and Medical Research Council, MRC Centre, Hills Road, Cambridge CB2 2QH, United Kingdom.

SOURCE: Proceedings of the National Academy of Sciences of the United States of America, (2002 Aug 20) 99 (17) 11085-90. Journal code: 7505876. ISSN: 0027-8424.

PUB. COUNTRY: United States

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

LANGUAGE: English

FILE SEGMENT: Priority Journals

ENTRY MONTH: 200209

ENTRY DATE: Entered STN: 20020822

Last Updated on STN: 20030105

Entered Medline: 20020927

TI CRINEPT-TROSY NMR reveals **p53** core domain bound in an unfolded form to the **chaperone** Hsp90.

AB The molecular **chaperone** Hsp90 sequesters oncogenic **mutants** of the tumor suppressor **p53** that have unstable core domains. It is not known whether **p53** is bound in an unfolded, partly folded, or distorted structure, as is unknown for the structure of any bound substrate. . . detail the structures of large complexes in which one component is (partly) unfolded. We have shown by transverse relaxation-optimized NMR **spectroscopy** combined with cross-correlated relaxation-enhanced polarization transfer (CRINEPT-TROSY) that **p53** core domain bound in an approximately 200-kDa complex with Hsp90 was predominantly unfolded lacking helical or sheet secondary structure. This. . .

CT Check Tags: Support, Non-U.S. Gov't
Binding Sites

DELACROIX

09/863,976

*Heat-Shock Proteins 90: CH, chemistry
Heat-Shock Proteins 90: ME, metabolism
 Magnetic Resonance Spectroscopy: MT, methods
Models, Molecular
 Protein Conformation
Protein Denaturation
Protein Folding
 *Protein p53: CH, chemistry
 Protein p53: ME, metabolism
Recombinant Proteins: CH, chemistry
Recombinant Proteins: ME, metabolism
Spectrometry, Fluorescence
Thermodynamics

CN 0 (Heat-Shock Proteins 90); 0 (Protein p53); 0 (Recombinant Proteins)

=>

09/863,976

(FILE 'HOME' ENTERED AT 22:06:34 ON 24 MAY 2004)

FILE 'HCAPLUS, BIOSIS, EMBASE, MEDLINE' ENTERED AT 22:06:56 ON 24 MAY 2004

L1 85 S P53 AND (CONFORMATION? OR STABILI?) AND CHAPERON? AND (MUTANT
L2 33 DUP REM L1 (52 DUPLICATES REMOVED)
L3 3 S L1 AND SPECTROSCOP?
L4 657 S (CONFORMATION? OR STABILI?) AND CHAPERON? AND SPECTROSCOP?
L5 326 DUP REM L4 (331 DUPLICATES REMOVED)
L6 64 S L5 AND (MUTANT? OR WILD(2A)TYPE?)
L7 12 S L6 AND (IDENTIFY? OR ASSAY?)

FILE 'STNGUIDE' ENTERED AT 22:19:23 ON 24 MAY 2004

FILE 'HCAPLUS, BIOSIS, EMBASE, MEDLINE' ENTERED AT 22:20:06 ON 24 MAY 2004

=> s l6 and py<=1998

3 FILES SEARCHED...

L8 14 L6 AND PY<=1998

=> d l8 abs ibib kwic 1-14

L8 ANSWER 1 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB The formation of active membrane-bound nitrate reductase A in Escherichia coli requires the presence of three subunits, NarG, NarH and NarI, as well as fourth protein, NarJ, that is not part of the active nitrate reductase. In narJ strains, both NarG and NarH subunits are associated in an unstable and inactive NarGH complex. A significant activation of this complex was observed in vitro after adding purified NarJ-6His polypeptide to the cell supernatant of a narJ strain. Once the apo-enzyme NarGHI of a narJ **mutant** has become anchored to the membrane via the NarI subunit, it cannot be reactivated by NarJ in vitro. NarJ protein specifically recognizes the catalytic NarG subunit. Fluorescence, ESR (EPR) **spectroscopy** and molybdenum quantification/based on inductively coupled plasma emission **spectroscopy** (ICPES) clearly indicate that, in the absence of NarJ, no molybdenum cofactor is present in the NarGH complex. We propose that NarJ is a specific **chaperone** that binds to NarG and may thus keep it in an appropriate competent-open **conformation** for the molybdenum cofactor insertion to occur, resulting in a catalytically active enzyme. Upon insertion of the molybdenum cofactor into the apo-nitrate reductase, NarJ is then dissociated from the activated enzyme.

ACCESSION NUMBER: 1998:319533 HCAPLUS

DOCUMENT NUMBER: 129:51404

TITLE: NarJ is a specific **chaperone** required for molybdenum cofactor assembly in nitrate reductase A of Escherichia coli

AUTHOR(S): Blasco, Francis; Dos Santos, Jean-Philippe; Magalon, Axel; Frixon, Chantal; Guigliarelli, Bruno; Santini, Claire-Lise; Giordano, Gerard

CORPORATE SOURCE: Laboratoire de Chimie Bacterienne, IBSM, CNRS, Marseille, 13402, Fr.

SOURCE: Molecular Microbiology (1998), 28(3), 435-447

CODEN: MOMIEE; ISSN: 0950-382X

PUBLISHER: Blackwell Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

DELACROIX

- TI NarJ is a specific **chaperone** required for molybdenum cofactor assembly in nitrate reductase A of Escherichia coli
- SO Molecular Microbiology (1998), 28(3), 435-447
CODEN: MOMIEE; ISSN: 0950-382X
- AB The formation of active membrane-bound nitrate reductase A in Escherichia coli requires the presence of three subunits, NarG, NarH and NarI, as well as fourth protein, NarJ, that is not part of the active nitrate reductase. In narJ strains, both NarG and NarH subunits are associated in an unstable and inactive NarGH complex. A significant activation of this complex was observed in vitro after adding purified NarJ-6His polypeptide to the cell supernatant of a narJ strain. Once the apo-enzyme NarGHI of a narJ mutant has become anchored to the membrane via the NarI subunit, it cannot be reactivated by NarJ in vitro. NarJ protein specifically recognizes the catalytic NarG subunit. Fluorescence, ESR (EPR) **spectroscopy** and molybdenum quantification/based on inductively coupled plasma emission **spectroscopy** (ICPES) clearly indicate that, in the absence of NarJ, no molybdenum cofactor is present in the NarGH complex. We propose that NarJ is a specific **chaperone** that binds to NarG and may thus keep it in an appropriate competent-open **conformation** for the molybdenum cofactor insertion to occur, resulting in a catalytically active enzyme. Upon insertion of the molybdenum cofactor into the apo-nitrate reductase, NarJ is then dissociated from the activated enzyme.
- ST NarJ **chaperone** molybdenum cofactor nitrate reductase;
Escherichia NarJ molybdenum cofactor nitrate reductase
- IT Escherichia coli
(NarJ subunit is a specific **chaperone** that binds to NarG subunit and is required for molybdenum cofactor assembly in nitrate reductase A of Escherichia coli)
- IT 9013-03-0, Nitrate reductase
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
(A; NarJ subunit is a specific **chaperone** that binds to NarG subunit and is required for molybdenum cofactor assembly in nitrate reductase A of Escherichia coli)
- IT 73508-07-3, Molybdoenzyme molybdenum cofactor
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(NarJ subunit is a specific **chaperone** that binds to NarG subunit and is required for molybdenum cofactor assembly in nitrate reductase A of Escherichia coli)
- L8 ANSWER 2 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN
- AB The subunit mol. mass of α -crystallin, like many small heat-shock proteins (sHsps), is around 20 kDa although the protein exists as a large aggregate of average mass around 800 kDa. Despite this large size, a well-resolved 1H NMR spectrum is observed for α -crystallin which arises from short, polar, highly-flexible and solvent-exposed C-terminal extensions in each of the subunits, α A- and α B-crystallin. These extensions are not involved in interactions with other proteins (e.g. β - And γ -crystallins) under non- **chaperone** conditions. As determined by NMR studies on **mutants** of α A-crystallin with alterations in its C-terminal extension, the extensions have an important role in acting as solubilizing agents for the relatively hydrophobic α -crystallin mol. and the high-mol.-weight (HMW) complex that forms during the **chaperone** action. The related sHsp, Hsp25, also exhibits a flexible C-terminal extension. Under **chaperone** conditions, and in the HMW complex isolated from old lenses, the C-terminal extension of the α A-crystallin subunit

maintains its flexibility whereas the α B-crystallin subunit loses, at least partially, its flexibility, implying that it is involved in interaction with the 'substrate' protein. The **conformation** of 'substrate' proteins when they interact with α -crystallin has been probed by ^1H NMR **spectroscopy** and it is concluded that α -crystallin interacts with 'substrate' proteins that are in a disordered molten globule state, but only when this state is on its way to large-scale aggregation and precipitation. By monitoring the ^1H and ^{31}P NMR spectra of α -crystallin in the presence of increasing concns. of urea, it is proposed that α -crystallin adopts a two-domain structure with the larger C-terminal domain unfolding first in the presence of denaturant. All these data have been combined into a model for the quaternary structure of α -crystallin. The model has two layers each of approx. 40 subunits arranged in an annulus or toroid. A large central cavity is present whose entrance is ringed by the flexible C-terminal extensions. A large hydrophobic region in the aggregate is exposed to solution and is available for interaction with 'substrate' proteins during the **chaperone** action.

ACCESSION NUMBER: 1998:264678 HCAPLUS
 DOCUMENT NUMBER: 129:64449
 TITLE: NMR **spectroscopy** of α -crystallin.
 Insights into the structure, interactions and
~~chaperone~~ action of small heat-shock proteins
 AUTHOR(S): Carver, John A.; Lindner, Robyn A.
 CORPORATE SOURCE: Department of Chemistry, The University of Wollongong,
 Wollongong, 2522, Australia
 SOURCE: International Journal of Biological Macromolecules (
 1998), 22(3,4), 197-209
 CODEN: IJBMDR; ISSN: 0141-8130
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 REFERENCE COUNT: 53 THERE ARE 53 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI NMR **spectroscopy** of α -crystallin. Insights into the
 structure, interactions and **chaperone** action of small heat-shock
 proteins
 SO International Journal of Biological Macromolecules (1998),
 22(3,4), 197-209
 CODEN: IJBMDR; ISSN: 0141-8130
 AB The subunit mol. mass of α -crystallin, like many small heat-shock
 proteins (sHsps), is around 20 kDa although the protein exists as a large
 aggregate of average mass around 800 kDa. Despite this large size, a
 well-resolved ^1H NMR spectrum is observed for α -crystallin which arises
 from short, polar, highly-flexible and solvent-exposed C-terminal
 extensions in each of the subunits, α A- and α B-crystallin.
 These extensions are not involved in interactions with other proteins
 (e.g. β - And γ -crystallins) under non- **chaperone**
 conditions. As determined by NMR studies on **mutants** of
 α A-crystallin with alterations in its C-terminal extension, the
 extensions have an important role in acting as solubilizing agents for the
 relatively hydrophobic α -crystallin mol. and the high-mol.-weight (HMW)
 complex that forms during the **chaperone** action. The related
 sHsp, Hsp25, also exhibits a flexible C-terminal extension. Under
chaperone conditions, and in the HMW complex isolated from old
 lenses, the C-terminal extension of the α A-crystallin subunit
 maintains its flexibility whereas the α B-crystallin subunit loses,
 at least partially, its flexibility, implying that it is involved in
 interaction with the 'substrate' protein. The **conformation** of

'substrate' proteins when they interact with α -crystallin has been probed by 1H NMR **spectroscopy** and it is concluded that α -crystallin interacts with 'substrate' proteins that are in a disordered molten globule state, but only when this state is on its way to large-scale aggregation and precipitation. By monitoring the 1H and 31P NMR spectra of α -crystallin in the presence of increasing concns. of urea, it is proposed that α -crystallin adopts a two-domain structure with the larger C-terminal domain unfolding first in the presence of denaturant. All these data have been combined into a model for the quaternary structure of α -crystallin. The model has two layers each of approx. 40 subunits arranged in an annulus or toroid. A large central cavity is present whose entrance is ringed by the flexible C-terminal extensions. A large hydrophobic region in the aggregate is exposed to solution and is available for interaction with 'substrate' proteins during the **chaperone** action.

- ST NMR **spectroscopy** alpha crystallin structure sHSP;
chaperone heat shock protein interaction crystallin; quaternary structure model alpha crystallin **conformation**
- IT Crystal structure
Hydrophobicity
Molecular modeling
NMR (nuclear magnetic resonance)
(NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT **Chaperonins**
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT Heat-shock proteins
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(low-mol.-weight; NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT Quaternary structure
(protein, model of; NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT **Conformation**
(protein; NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT Crystallins
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(α -; NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT Crystallins
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(α A-; NMR **spectroscopy** of α -crystallin: structure, interactions and **chaperone** action of small heat-shock proteins)
- IT Crystallins
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP

(Properties); BIOL (Biological study); PROC (Process)
 (α B-; NMR **spectroscopy** of α -crystallin:
 structure, interactions and **chaperone** action of small
 heat-shock proteins)

L8 ANSWER 3 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB Defects in electron transfer flavoprotein (ETF) or its electron acceptor, electron transfer flavoprotein-ubiquinone oxidoreductase (ETF-QO), cause the human inherited metabolic disease glutaric acidemia type II. In this disease, electron transfer from nine primary flavoprotein dehydrogenases to the main respiratory chain is impaired. Among these dehydrogenases are the four chain length-specific flavoprotein dehydrogenases of fatty acid β -oxidation. In this investigation, two mutations in the α subunit that have been identified in patients were expressed in *Escherichia coli*. Of the two **mutant** alleles, α T266M and α G116R, the former is the most frequent mutation found in patients with ETF deficiency. The crystal structure of human ETF shows that α G116 lies in a hydrophobic pocket, under a contact residue of the α/β subunit interface, and that the hydroxyl hydrogen of α T266 is hydrogen-bonded to N(5) of the FAD; the amide backbone hydrogen of α T266 is hydrogen-bonded to C(4)-O of the flavin prosthetic group (Roberts, D. L., Frerman, F. E. and Kim, J-J. P. (1996) Proc. Natl. Acad. Sci. U. S. A. 93, 14355-14360). Stable expression of the α G116R ETF required coexpression of the **chaperonins**, GroEL and GroES. α G116R ETF folds into a **conformation** different from the **wild type**, and is catalytically inactive in crude exts. It is unstable and could not be extensively purified. The α T266M ETF was purified and characterized after **stabilization** to proteolysis in crude exts. Although the global structure of this **mutant** protein is unchanged, its flavin environment is altered as indicated by absorption and CD **spectroscopy** and the kinetics of flavin release from the oxidized and reduced protein. The loss of the hydrogen bond at N(5) of the flavin and the altered flavin binding increase the thermodyn. **stability** of the flavin semiquinone by 10-fold relative to the semiquinone of **wild type** ETF. The mutation has relatively little effect on the reductive half-reaction of ETF catalyzed by sarcosine and medium chain acyl-CoA dehydrogenases which reduce the flavin to the semiquinone. However, k_{cat}/K_m of ETF-QO in a coupled acyl-CoA:ubiquinone reductase assay with oxidized α T266M ETF as substrate is reduced 33-fold; this decrease is due in largest part to a decrease in the rate of disproportionation of the α T266M ETF semiquinone catalyzed by ETF-QO.

ACCESSION NUMBER: 1997:689123 HCAPLUS
 DOCUMENT NUMBER: 128:1285
 TITLE: Expression and characterization of two pathogenic mutations in human electron transfer flavoprotein
 AUTHOR(S): Salazar, Denise; Zhang, Lening; Degala, Gregory D.; Frerman, Frank E.
 CORPORATE SOURCE: Program in Cellular and Developmental Biology and the Department of Pediatrics, University of Colorado School of Medicine, Denver, CO, 80262, USA
 SOURCE: Journal of Biological Chemistry (1997), 272(42), 26425-26433
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular Biology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

- SO Journal of Biological Chemistry (1997), 272(42), 26425-26433
CODEN: JBCHA3; ISSN: 0021-9258
- AB Defects in electron transfer flavoprotein (ETF) or its electron acceptor, electron transfer flavoprotein-ubiquinone oxidoreductase (ETF-QO), cause the human inherited metabolic disease glutaric acidemia type II. In this disease, electron transfer from nine primary flavoprotein dehydrogenases to the main respiratory chain is impaired. Among these dehydrogenases are the four chain length-specific flavoprotein dehydrogenases of fatty acid β -oxidation. In this investigation, two mutations in the α subunit that have been identified in patients were expressed in *Escherichia coli*. Of the two **mutant** alleles, α T266M and α G116R, the former is the most frequent mutation found in patients with ETF deficiency. The crystal structure of human ETF shows that α G116 lies in a hydrophobic pocket, under a contact residue of the α/β subunit interface, and that the hydroxyl hydrogen of α T266 is hydrogen-bonded to N(5) of the FAD; the amide backbone hydrogen of α T266 is hydrogen-bonded to C(4)-O of the flavin prosthetic group (Roberts, D. L., Frerman, F. E. and Kim, J.-J. P. (1996) *Proc. Natl. Acad. Sci. U. S. A.* 93, 14355-14360). Stable expression of the α G116R ETF required coexpression of the **chaperonins**, GroEL and GroES. α G116R ETF folds into a **conformation** different from the **wild type**, and is catalytically inactive in crude exts. It is unstable and could not be extensively purified. The α T266M ETF was purified and characterized after **stabilization** to proteolysis in crude exts. Although the global structure of this **mutant** protein is unchanged, its flavin environment is altered as indicated by absorption and CD **spectroscopy** and the kinetics of flavin release from the oxidized and reduced protein. The loss of the hydrogen bond at N(5) of the flavin and the altered flavin binding increase the thermodyn. **stability** of the flavin semiquinone by 10-fold relative to the semiquinone of **wild type** ETF. The mutation has relatively little effect on the reductive half-reaction of ETF catalyzed by sarcosine and medium chain acyl-CoA dehydrogenases which reduce the flavin to the semiquinone. However, k_{cat}/K_m of ETF-QO in a coupled acyl-CoA:ubiquinone reductase assay with oxidized α T266M ETF as substrate is reduced 33-fold; this decrease is due in largest part to a decrease in the rate of disproportionation of the α T266M ETF semiquinone catalyzed by ETF-QO.
- IT **Chaperonins**
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(GroEL; expression and characterization of two pathogenic mutations in human electron transfer flavoprotein)
- IT **Chaperonins**
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(GroES; expression and characterization of two pathogenic mutations in human electron transfer flavoprotein)
- IT **Conformation**
(protein; expression and characterization of two pathogenic mutations in human electron transfer flavoprotein)
- L8 ANSWER 4 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN
- AB α -Crystallins occur as multimeric complexes, which are able to suppress precipitation of unfolding proteins. Although the mechanism of this **chaperone**-like activity is unknown, the affinity of

α -crystallin for aggregation-prone proteins is probably based on hydrophobic interactions. α -Crystallins expose a considerable hydrophobic surface to solution, but nevertheless they are very stable and highly soluble. An explanation for this paradox may be that α -crystallin subunits have a polar and unstructured C-terminal extension that functions as a sort of solubilizer. In this paper we have described five α A-crystallins in which charged and hydrophobic residues were inserted in the C-terminal extension. Introduction of lysine, arginine, and aspartate does not substantially influence **chaperone**-like activity. In contrast, introduction of a hydrophobic tryptophan greatly diminishes functional activity. CD expts. indicate that this **mutant** has a normal secondary structure and fluorescence measurements show that the inserted tryptophan is located in polar environment. However, NMR **spectroscopy** clearly demonstrates that the presence of the tryptophan residue dramatically reduces the flexibility of the C-terminal extension. Furthermore, the introduction of this tryptophan results in a considerably decreased thermostability of the protein. We conclude that changing the polarity of the C-terminal extension of α A-crystallin by insertion of a highly hydrophobic residue can seriously disturb structural and functional integrity.

ACCESSION NUMBER: 1996:716584 HCAPLUS
 DOCUMENT NUMBER: 126:28206
 TITLE: Immobilization of the C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity
 AUTHOR(S): Smulders, Ronald H. P. H.; Carver, John A.; Lindner, Robyn A.; van Boeckel, Martinus A. M.; Bloemendal, Hans; de Jong, Wilfried W.
 CORPORATE SOURCE: Dep. Biochem., Univ Nijmegen, Nijmegen, 6500 HB, Neth.
 SOURCE: Journal of Biological Chemistry (1996), 271(46), 29060-29066
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular Biology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 TI Immobilization of the C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity
 SO Journal of Biological Chemistry (1996), 271(46), 29060-29066
 CODEN: JBCHA3; ISSN: 0021-9258
 AB α -Crystallins occur as multimeric complexes, which are able to suppress precipitation of unfolding proteins. Although the mechanism of this **chaperone**-like activity is unknown, the affinity of α -crystallin for aggregation-prone proteins is probably based on hydrophobic interactions. α -Crystallins expose a considerable hydrophobic surface to solution, but nevertheless they are very stable and highly soluble. An explanation for this paradox may be that α -crystallin subunits have a polar and unstructured C-terminal extension that functions as a sort of solubilizer. In this paper we have described five α A-crystallins in which charged and hydrophobic residues were inserted in the C-terminal extension. Introduction of lysine, arginine, and aspartate does not substantially influence **chaperone**-like activity. In contrast, introduction of a hydrophobic tryptophan greatly diminishes functional activity. CD expts. indicate that this **mutant** has a normal secondary structure and fluorescence measurements show that the inserted tryptophan is located in polar environment. However, NMR **spectroscopy** clearly demonstrates that the presence of the tryptophan residue dramatically

reduces the flexibility of the C-terminal extension. Furthermore, the introduction of this tryptophan results in a considerably decreased thermostability of the protein. We conclude that changing the polarity of the C-terminal extension of α A-crystallin by insertion of a highly hydrophobic residue can seriously disturb structural and functional integrity.

ST crystallin structure **chaperone** function **conformation**

IT **Conformation**

(immobilization of C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity)

IT **Chaperonins**

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process)

(immobilization of C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity)

IT Crystallins

RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process)

(α A-; immobilization of C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity)

IT 73-22-3, Tryptophan, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(immobilization of C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity)

IT 9004-10-8, Insulin, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); PEP (Physical, engineering or chemical process); BIOL (Biological study); PROC (Process)

(immobilization of C-terminal extension of bovine α A-crystallin reduces **chaperone**-like activity)

L8 ANSWER 5 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB α -Crystallin, the major protein of the ocular lens, acts as a mol. **chaperone** by suppressing the nonspecific aggregation of damaged proteins. To investigate the mechanism of the interaction between α -crystallin and substrate proteins, we prepared a tryptophan-free **mutant** of human α A-crystallin and assessed the **conformation** of thermally destabilized proteins captured by this **chaperone** using fluorescence **spectroscopy**. The fluorescence emission characteristics of bound substrates (rhodanese and γ -crystallin) and the results of fluorescence quenching expts. indicate that the proteins captured by α -crystallin are characterized by a very low degree of unfolding. In particular, the structure of rhodanese bound to α A-crystallin appears to be considerably more native-like compared to that of the enzyme bound to the **chaperonin** GroEL. We postulate that α -crystallin (and likely other small heat shock proteins) recognize preferentially the aggregation-prone conformers that occur very early on the denaturation pathway. With its ability to capture and **stabilize** these early non-native structures, α -crystallin appears to be uniquely well suited to **chaperone** the transparency properties of the ocular lens.

ACCESSION NUMBER: 1996:282924 HCAPLUS

DOCUMENT NUMBER: 124:335953

TITLE: **Conformational** properties of substrate proteins bound to a molecular **chaperone**

AUTHOR(S): α -crystallin
 CORPORATE SOURCE: Das, Kali P.; Petrash, J. Mark; Surewicz, Witold K.
 SOURCE: Mason Eye Inst and Dep. Biochem., Univ. Missouri,
 Columbia, MO, 65212, USA
 PUBLISHER: Journal of Biological Chemistry (1996),
 271(18), 10449-10452
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular
 Biology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

TI **Conformational** properties of substrate proteins bound to a
 molecular **chaperone** α -crystallin
 SO Journal of Biological Chemistry (1996), 271(18), 10449-10452
 CODEN: JBCHA3; ISSN: 0021-9258
 AB α -Crystallin, the major protein of the ocular lens, acts as a mol.
chaperone by suppressing the nonspecific aggregation of damaged
 proteins. To investigate the mechanism of the interaction between
 α -crystallin and substrate proteins, we prepared a tryptophan-free
mutant of human α A-crystallin and assessed the
conformation of thermally destabilized proteins captured by this
chaperone using fluorescence **spectroscopy**. The
 fluorescence emission characteristics of bound substrates (rhodanese and
 γ -crystallin) and the results of fluorescence quenching expts.
 indicate that the proteins captured by α -crystallin are
 characterized by a very low degree of unfolding. In particular, the
 structure of rhodanese bound to α A-crystallin appears to be
 considerably more native-like compared to that of the enzyme bound to the
chaperonin GroEL. We postulate that α -crystallin (and
 likely other small heat shock proteins) recognize preferentially the
 aggregation-prone conformers that occur very early on the denaturation
 pathway. With its ability to capture and **stabilize** these early
 non-native structures, α -crystallin appears to be uniquely well
 suited to **chaperone** the transparency properties of the ocular
 lens.
 ST crystallin **chaperone** protein **conformation**
 IT **Conformation** and Conformers
 (conformational properties of substrate proteins bound to
 mol. **chaperone** α -crystallin)
 IT Proteins, biological studies
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP
 (Properties); BIOL (Biological study); PROC (Process)
 (conformational properties of substrate proteins bound to
 mol. **chaperone** α -crystallin)
 IT Crystallins
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (α A-, **conformational** properties of substrate proteins
 bound to mol. **chaperone** α -crystallin)
 IT Crystallins
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP
 (Properties); BIOL (Biological study); PROC (Process)
 (β L-, **conformational** properties of substrate proteins
 bound to mol. **chaperone** α -crystallin)
 IT Crystallins
 RL: BPR (Biological process); BSU (Biological study, unclassified); PRP
 (Properties); BIOL (Biological study); PROC (Process)
 (γ -, **conformational** properties of substrate proteins
 bound to mol. **chaperone** α -crystallin)

IT 9026-04-4, Rhodanese

RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)
(**conformational** properties of substrate proteins bound to mol. **chaperone** α -crystallin)

L8 ANSWER 6 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB A review, with .apprx.103 refs. The molten globule state of α -lactalbumin is the best-characterized folding intermediate of globular proteins and has been studied intensively by various **spectroscopic** and physicochem. techniques, including stopped-flow CD and fluorescence **spectroscopies**, a hydrogen-exchange technique, $^1\text{H-NMR}$ **spectroscopy**, disulfide-exchange chemical, site-directed mutagenesis, and calorimetric techniques. This review summarizes recent studies. Major findings about the structure of the molten globule state are: (1) it is highly heterogeneous, having a highly structured α -helical domain with the β -sheet domain being significantly unfolded; and (2) it is not a nonspecific, collapsed polypeptide but already has a native-like tertiary fold. These structural characteristics are essential to fully understand the thermodyn. properties of the molten globule state, which are described in connection with a recently proposed computational approach to predict the structure of the molten globule state of a protein. **Mutant** proteins in which the **stability** of the molten globule state was changed were constructed. Studies of the equilibrium unfolding and kinetic refolding of the **mutant** proteins will provide further insight into the molten globule state as a folding intermediate. In spite of an initial expectation that the structure recognized by an Escherichia coli **chaperone**, GroEL, is the molten globule, the interaction of GroEL with α -lactalbumin in the molten globule state is much weaker than the interaction with more unfolded states of α -lactalbumin, a disulfide-reduced form, and disulfide rearranged species.

ACCESSION NUMBER: 1996:78612 HCAPLUS
DOCUMENT NUMBER: 124:109868
TITLE: The molten globule state of α -lactalbumin
AUTHOR(S): Kuwajima, Kunihiro
CORPORATE SOURCE: Sch. Sci., Univ. Tokyo, Tokyo, 113, Japan
SOURCE: FASEB Journal (1996), 10(1), 102-9
CODEN: FAJOEC; ISSN: 0892-6638
PUBLISHER: Federation of American Societies for Experimental Biology
DOCUMENT TYPE: Journal; General Review
LANGUAGE: English

SO FASEB Journal (1996), 10(1), 102-9
CODEN: FAJOEC; ISSN: 0892-6638

AB A review, with .apprx.103 refs. The molten globule state of α -lactalbumin is the best-characterized folding intermediate of globular proteins and has been studied intensively by various **spectroscopic** and physicochem. techniques, including stopped-flow CD and fluorescence **spectroscopies**, a hydrogen-exchange technique, $^1\text{H-NMR}$ **spectroscopy**, disulfide-exchange chemical, site-directed mutagenesis, and calorimetric techniques. This review summarizes recent studies. Major findings about the structure of the molten globule state are: (1) it is highly heterogeneous, having a highly structured α -helical domain with the β -sheet domain being significantly unfolded; and (2) it is not a nonspecific, collapsed polypeptide but already has a native-like tertiary fold. These structural characteristics are essential to fully understand the thermodyn. properties of the molten globule state, which are described in connection with a

recently proposed computational approach to predict the structure of the molten globule state of a protein. **Mutant** proteins in which the **stability** of the molten globule state was changed were constructed. Studies of the equilibrium unfolding and kinetic refolding of the **mutant** proteins will provide further insight into the molten globule state as a folding intermediate. In spite of an initial expectation that the structure recognized by an Escherichia coli **chaperone**, GroEL, is the molten globule, the interaction of GroEL with α -lactalbumin in the molten globule state is much weaker than the interaction with more unfolded states of α -lactalbumin, a disulfide-reduced form, and disulfide rearranged species.

ST review lactalbumin **conformation** molten globule

IT **Conformation** and Conformers

(molten globule; molten globule state of α -lactalbumin)

L8 ANSWER 7 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB Temperature-sensitive folding (tsf) **mutants** of the phage P22 coat protein prevent newly synthesized polypeptide chains from reaching the **conformation** competent for capsid assembly in cells and can be rescued by the GroEL **chaperone**. Here, the **stabilities** of **wild-type** and 4 tsf **mutant** unpolymd. subunits were investigated. **Wild-type** coat protein subunits denatured at 40°, with a calorimetric enthalpy of .apprx.600 kJ/mol. Comparison with coat protein denaturation within the shell lattice ($T_m = 87^\circ$, ΔH .apprx. 1700 kJ/mol) indicates that protein-protein interactions within the capsid provide enormous **stabilization**. The melting temps. of the subunits carrying tsf substitutions were similar to **wild-type**. At low temps., the tsf **mutants**, but not the **wild-type**, formed non-covalent dimers, which were dissociated at temps. $>30^\circ$. **Spectroscopic** and calorimetric studies indicated that the **mutant** proteins have reduced amts. of ordered structure at low temperature, as compared to the **wild-type** protein. Although complex, the in vitro phenotypes are consistent with the in vivo finding that the **mutants** are defective in folding, rather than subunit **stability**. These results suggest a role for incompletely folded subunits as precursors in viral capsid assembly, providing a mechanism of reaching multiple **conformations** in the polymerized form.

ACCESSION NUMBER: 1995:697734 HCAPLUS

DOCUMENT NUMBER: 123:79168

TITLE: **Stability of wild-type**
and temperature-sensitive protein subunits of the
phage P22 capsid

AUTHOR(S): Galisteo, Maria L.; Gordon, Carl L.; King, Jonathan

CORPORATE SOURCE: Fac. Cienc., Univ. Granada, Granada, 18071, Spain

SOURCE: Journal of Biological Chemistry (1995),
270(28), 16595-601

CODEN: JBCHA3; ISSN: 0021-9258

PUBLISHER: American Society for Biochemistry and Molecular Bio
logy

DOCUMENT TYPE: Journal

LANGUAGE: English

TI **Stability of wild-type** and
temperature-sensitive protein subunits of the phage P22 capsid
SO Journal of Biological Chemistry (1995), 270(28), 16595-601
CODEN: JBCHA3; ISSN: 0021-9258

AB Temperature-sensitive folding (tsf) **mutants** of the phage P22 coat protein prevent newly synthesized polypeptide chains from reaching the **conformation** competent for capsid assembly in cells and can be

rescued by the GroEL **chaperone**. Here, the **stabilities** of **wild-type** and 4 tsf mutant unpolymd. subunits were investigated. **Wild-type** coat protein subunits denatured at 40°, with a calorimetric enthalpy of .apprx.600 kJ/mol. Comparison with coat protein denaturation within the shell lattice ($T_m = 87^\circ$, ΔH .apprx. 1700 kJ/mol) indicates that protein-protein interactions within the capsid provide enormous **stabilization**. The melting temps. of the subunits carrying tsf substitutions were similar to **wild-type**. At low temps., the tsf **mutants**, but not the **wild-type**, formed non-covalent dimers, which were dissociated at temps. $>30^\circ$. **Spectroscopic** and calorimetric studies indicated that the **mutant** proteins have reduced amts. of ordered structure at low temperature, as compared to the **wild-type** protein. Although complex, the in vitro phenotypes are consistent with the in vivo finding that the **mutants** are defective in folding, rather than subunit **stability**. These results suggest a role for incompletely folded subunits as precursors in viral capsid assembly, providing a mechanism of reaching multiple **conformations** in the polymerized form.

ST protein **stability** phage P22 capsid

IT Virus, bacterial

(P22, **stability** of **wild-type** and

temperature-sensitive protein subunits of the phage P22 capsid)

IT Proteins, specific or class

RL: PRP (Properties)

(capsid, **stability** of **wild-type** and

temperature-sensitive protein subunits of the phage P22 capsid)

L8 ANSWER 8 OF 14 HCAPLUS COPYRIGHT 2004 ACS on STN

AB Three **mutants** of barnase and a pro-barnase variant, which have a variety of different phys. properties but the same overall protein structure, were analyzed for their folding in the presence of the mol. **chaperone** GroEL. **Mutants** were chosen on the basis that changes in their refolding rate consts. in solution are not correlated with the changes in their **stability**. All barnase variants fold considerably more slowly when bound to GroEL. However, barnase refolding on GroEL parallels that in solution: there is a linear relationship between the refolding rate consts., obtained for **wild-type** and all **mutants** of barnase, in the presence and absence of GroEL. Barnase is synthesized in vivo with a 13 amino acid pro-sequence attached to the N-terminus. The pro-sequence of pro-barnase is shown by NMR **spectroscopy** to be devoid of defined structure. The presence of this pro-sequence has no effect on the overall refolding rate constant or the activity of barnase. In the presence of GroEL, the refolding of pro-barnase is retarded relatively more strongly than that of **wild-type** and the **mutant** barnase proteins, suggesting that the pro-sequence provides addnl. binding sites for the **chaperone**

ACCESSION NUMBER: 1994:100398 HCAPLUS

DOCUMENT NUMBER: 120:100398

TITLE: Refolding of barnase **mutants** and pro-barnase in the presence and absence of GroEL

AUTHOR(S): Gray, Tamara E.; Eder, Joerg; Bycroft, Mark; Day, Anthony G.; Fersht, Alan R.

CORPORATE SOURCE: Chem. Lab., Univ. Cambridge, Cambridge, CB2 1EW, UK

SOURCE: EMBO Journal (1993), 12(11), 4145-50

CODEN: EMJODG; ISSN: 0261-4189

DOCUMENT TYPE: Journal

LANGUAGE: English

- TI Refolding of barnase **mutants** and pro-barnase in the presence and absence of GroEL
- SO EMBO Journal (1993), 12(11), 4145-50
CODEN: EMJODG; ISSN: 0261-4189
- AB Three **mutants** of barnase and a pro-barnase variant, which have a variety of different phys. properties but the same overall protein structure, were analyzed for their folding in the presence of the mol. **chaperone** GroEL. **Mutants** were chosen on the basis that changes in their refolding rate consts. in solution are not correlated with the changes in their **stability**. All barnase variants fold considerably more slowly when bound to GroEL. However, barnase refolding on GroEL parallels that in solution: there is a linear relationship between the refolding rate consts., obtained for **wild-type** and all **mutants** of barnase, in the presence and absence of GroEL. Barnase is synthesized in vivo with a 13 amino acid pro-sequence attached to the N-terminus. The pro-sequence of pro-barnase is shown by NMR **spectroscopy** to be devoid of defined structure. The presence of this pro-sequence has no effect on the overall refolding rate constant or the activity of barnase. In the presence of GroEL, the refolding of pro-barnase is retarded relatively more strongly than that of **wild-type** and the **mutant** barnase proteins, suggesting that the pro-sequence provides addnl. binding sites for the **chaperone**
- ST **chaperone** GroEL barnase probarnase refolding; propeptide
chaperone GroEL interaction barnase refolding
- IT **Conformation** and Conformers
(of barnase and probarnase, refolding of, GroEL protein effect on)
- IT Proteins, specific or class
RL: BIOL (Biological study)
(**chaperonins** 60, refolding of barnase and probarnase in presence on)
- IT **Conformation** and Conformers
(random-coil, of barnase propeptide)
- IT 9026-12-4, Barnase
RL: BIOL (Biological study)
(refolding of **mutant** and native forms of, GroEL protein effect on)
- L8 ANSWER 9 OF 14 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
- AB The expression of the melanin operon (melC) of Streptomyces antibioticus requires the **chaperone**-like protein MelC1 for the incorporation of two copper ions (designated as CuA and CuB) and the secretion of the apotyrosinase (MelC2) via a transient binary complex formation between these two proteins. To investigate whether the copper ligand of tyrosinase is involved in this MelC1ntdotMelC2 binary complex function, six single substitution mutations were introduced into the CuA and CuB sites. These mutations led to differential effects on the **stability**, copper content, and export function of binary complexes but a complete abolishment of tyrosinase activity. The defects in the tyrosinase activity in **mutants** were not because of the impairment of the formation of MelC1ntdotMelC2 complex but rather the failure of MelC2 to be discharged from the copper-activated binary complex. Moreover, the impairments on the discharge of the **mutant** MelC2 from all the **mutant** binary complexes appeared to result from the structural changes in their apoforms or copper-activated forms of the complexes, as evidenced by the fluorescence emission and circular dichroism spectral analysis. Therefore, each of six copper ligands in Streptomyces tyrosinase binuclear copper sites plays a pivotal role in the final maturation and the discharge of tyrosinase from the binary complex

but has a less significant role in its secretion.

ACCESSION NUMBER: 1998:390187 BIOSIS

DOCUMENT NUMBER: PREV199800390187

TITLE: Roles of copper ligands in the activation and secretion of *Streptomyces tyrosinase*.

AUTHOR(S): Tsai, Tzung-Yuan; Lee, Yan-Hwa Wu [Reprint author]

CORPORATE SOURCE: Inst. Biochemistry, National Yang-Ming Univ., Taipei 112, Taiwan

SOURCE: Journal of Biological Chemistry, (July 24, 1998) Vol. 273, No. 30, pp. 19243-19250. print.
CODEN: JBCHA3. ISSN: 0021-9258.

DOCUMENT TYPE: Article

LANGUAGE: English

ENTRY DATE: Entered STN: 10 Sep 1998

Last Updated on STN: 21 Oct 1998

SO Journal of Biological Chemistry, (July 24, 1998) Vol. 273, No. 30, pp. 19243-19250. print.

CODEN: JBCHA3. ISSN: 0021-9258.

AB The expression of the melanin operon (*melC*) of *Streptomyces antibioticus* requires the **chaperone**-like protein *MelC1* for the incorporation of two copper ions (designated as *CuA* and *CuB*) and the secretion of the apotyrosinase. . . six single substitution mutations were introduced into the *CuA* and *CuB* sites. These mutations led to differential effects on the **stability**, copper content, and export function of binary complexes but a complete abolishment of tyrosinase activity. The defects in the tyrosinase activity in **mutants** were not because of the impairment of the formation of *MelC1*ntd*MelC2* complex but rather the failure of *MelC2* to be discharged from the copper-activated binary complex. Moreover, the impairments on the discharge of the **mutant** *MelC2* from all the **mutant** binary complexes appeared to result from the structural changes in their apoforms or copper-activated forms of the complexes, as evidenced. . .

IT Methods & Equipment

circular dichroism spectral analysis: analytical method,

spectroscopic techniques: CT; fluorescence emission

spectroscopy: analytical method, **spectroscopic**

techniques: CB; immunoaffinity chromatography: affinity chromatography,

analytical method; immunoblot analysis: Analysis/Characterization

Techniques: CB, analytical method; AVIV 60DS spectropolarimeter: AVIV

Associates, . . .

L8 ANSWER 10 OF 14 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

AB *DnaJ* is a molecular **chaperone**, which not only binds to its various protein substrates, but can also activate the *DnaK* co-**chaperone** to bind to its various protein substrates as well. *DnaJ* is a modular protein, which contains a putative zinc finger motif of unknown function. Quantitation of the released *Zn(II)* ions, upon challenge with p-hydroxymercuriphenylsulfonic acid, and by atomic absorption showed that two *Zn(II)* ions interact with each monomer of *DnaJ*. Following the release of *Zn(II)* ions, the free cysteine residues probably form disulfide bridge(s), which contribute to overcoming the destabilizing effect of losing *Zn(II)*. Supporting this view, infrared and circular dichroism studies show that the *DnaJ* secondary structure is largely unaffected by the release of *Zn(II)*. Moreover, infrared spectra recorded at different temperatures, as well as scanning calorimetry, show that the *Zn(H)* ions help to **stabilize** *DnaJ*'s tertiary structure. An internal 57-amino acid deletion of the cysteine-rich region did not noticeably affect the affinity of this **mutant** protein, *DnaJ-DELTA-144-200*, to bind *DnaK* nor its ability to stimulate *DnaK*'s

ATPase activity. However, the DnaJ-DELTA-144-200 was unable to induce DnaK to a **conformation** required for the **stabilization** of the DnaK-substrate complex. Additionally, the DnaJ-DELTA-144-200 **mutant** protein alone was unimpaired in its ability to interact with its sigma-32 transcription factor substrate, but exhibited reduced affinity toward its P1 RepA and lambda-P substrates. Finally, these in vitro results correlate well with the in vivo observed partial inhibition of bacteriophage lambda growth in a DnaJ-DELTA-144-200 **mutant** background.

ACCESSION NUMBER: 1996:327422 BIOSIS
DOCUMENT NUMBER: PREV199699049778
TITLE: Structure-function analysis of the zinc finger region of the DnaJ molecular **chaperone**.
AUTHOR(S): Banecki, Bogdan; Liberek, Krzysztof; Wall, Daniel; Wawrzynow, Alicja; Georgopoulos, Costa; Bertoli, Enrico; Tanfani, Fabio; Zylicz, Maciej [Reprint author]
CORPORATE SOURCE: Div. Biophys., Dep. Mol. Biol., Univ. Gdansk, 80-822 Gdansk, Kladki 24, Poland
SOURCE: Journal of Biological Chemistry, (1996) Vol. 271, No. 25, pp. 14840-14848.
CODEN: JBCHA3. ISSN: 0021-9258.
DOCUMENT TYPE: Article
LANGUAGE: English
ENTRY DATE: Entered STN: 26 Jul 1996
Last Updated on STN: 27 Jul 1996
TI Structure-function analysis of the zinc finger region of the DnaJ molecular **chaperone**.
SO Journal of Biological Chemistry, (1996) Vol. 271, No. 25, pp. 14840-14848.
CODEN: JBCHA3. ISSN: 0021-9258.
AB DnaJ is a molecular **chaperone**, which not only binds to its various protein substrates, but can also activate the DnaK co-**chaperone** to bind to its various protein substrates as well. DnaJ is a modular protein, which contains a putative zinc finger. . . Zn(II). Moreover, infrared spectra recorded at different temperatures, as well as scanning calorimetry, show that the Zn(H) ions help to **stabilize** DnaJ's tertiary structure. An internal 57-amino acid deletion of the cysteine-rich region did not noticeably affect the affinity of this **mutant** protein, DnaJ-DELTA-144-200, to bind DnaK nor its ability to stimulate DnaK's ATPase activity. However, the DnaJ-DELTA-144-200 was unable to induce DnaK to a **conformation** required for the **stabilization** of the DnaK-substrate complex. Additionally, the DnaJ-DELTA-144-200 **mutant** protein alone was unimpaired in its ability to interact with its sigma-32 transcription factor substrate, but exhibited reduced affinity toward. . . these in vitro results correlate well with the in vivo observed partial inhibition of bacteriophage lambda growth in a DnaJ-DELTA-144-200 **mutant** background.
IT Miscellaneous Descriptors
ANALYTICAL METHOD; BIOCHEMISTRY AND MOLECULAR BIOPHYSICS; CIRCULAR DICHROISM; DNAJ; IR **SPECTROSCOPY**; MOLECULAR **CHAPERONE**; P-HYDROXYPHENYLMERCURIC ACID; SIGMA FACTOR-32; STRUCTURE-FUNCTION ANALYSIS; TRANSCRIPTION FACTOR; TRANSCRIPTION FACTOR INTERACTION; ZINC FINGER REGION; ZINC ION
L8 ANSWER 11 OF 14 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
AB Methanol dehydrogenase (MDH) from Methylobacterium extorquens, Methylophilus methylotrophus. Paracoccus denitrificans and Hyphomicrobium X all contained a single atom of Ca²⁺ per $\alpha_2\beta_2$ tetramer. The

role of Ca^{2+} was investigated using the MDH from *Methylobacterium extorquens*. This was shown to be similar to the MDH from *Hyphomicrobium X* in having 2 mol of prosthetic group (pyrroloquinoline quinine; PQQ) per mol of tetramer, the PQQ being predominantly in the semiquinone form. MDH isolated from the methanol oxidation **mutants** MoxA-, K- and L- contained no Ca^{2+} . They were identical with the enzyme isolated from **wild-type** bacteria with respect to molecular size, subunit configuration, pI, N-terminal amino acid sequence and **stability** under denaturing conditions (low pH, high urea and high guanidinium chloride) and in the nature and content of the prosthetic group (2 mol of PQQ per mol of MDH). They differed in their lack of Ca^{2+} , the oxidation state of the extracted PQQ (fully oxidized), absence of the semiquinone form of PQQ in the enzyme, reactivity with the suicide inhibitor cyclopropanol and absorption spectrum, which indicated that PQQ is bound differently from that in normal MDH. Incubation of MDH from the **mutants** in calcium salts led to irreversible time-dependent reconstitution of full activity concomitant with restoration of a spectrum corresponding to that of fully reduced normal MDH. It is concluded that Ca^{2+} in MDH is directly or indirectly involved in binding PQQ in the active site. The MoxA-, K- and L- proteins may be involved in maintaining a high Ca^{2+} concentration in the periplasm. It is more likely, however, that they fill a '**chaperone**' function, **stabilizing** a configuration of MDH which permits incorporation of low concentrations of Ca^{2+} into the protein.

ACCESSION NUMBER: 92336416 EMBASE

DOCUMENT NUMBER: 1992336416

TITLE: Characterization of **mutant** forms of the quinoprotein methanol dehydrogenase lacking an essential calcium ion.

AUTHOR: Richardson I.W.; Anthony C.

CORPORATE SOURCE: SERC, Centre Molecular Recognition, Department of Biochemistry, University of Southampton, Southampton SO9 3TU, United Kingdom

SOURCE: Biochemical Journal, (1992) 287/3 (709-715).

ISSN: 0264-6021 CODEN: BIJOAK

COUNTRY: United Kingdom

DOCUMENT TYPE: Journal; Article

FILE SEGMENT: 004 Microbiology

029 Clinical Biochemistry

LANGUAGE: English

SUMMARY LANGUAGE: English

TI Characterization of **mutant** forms of the quinoprotein methanol dehydrogenase lacking an essential calcium ion.

SO Biochemical Journal, (1992) 287/3 (709-715).

ISSN: 0264-6021 CODEN: BIJOAK

AB . . . quinine; PQQ) per mol of tetramer, the PQQ being predominantly in the semiquinone form. MDH isolated from the methanol oxidation **mutants** MoxA-, K- and L- contained no Ca^{2+} . They were identical with the enzyme isolated from **wild-type** bacteria with respect to molecular size, subunit configuration, pI, N-terminal amino acid sequence and **stability** under denaturing conditions (low pH, high urea and high guanidinium chloride) and in the nature and content of the prosthetic. . . and absorption spectrum, which indicated that PQQ is bound differently from that in normal MDH. Incubation of MDH from the **mutants** in calcium salts led to irreversible time-dependent reconstitution of full activity concomitant with restoration of a spectrum corresponding to that. . . be involved in maintaining a high Ca^{2+} concentration in the periplasm. It is more likely, however, that they fill a '**chaperone**' function, **stabilizing** a configuration of

MDH which permits incorporation of low concentrations of Ca^{2+} into the protein.

CT Medical Descriptors:

***bacterium mutant**
 *enzyme structure
absorption spectroscopy
 amino acid sequence
 amino terminal sequence
 article
 controlled study
 enzyme active site
 enzyme analysis
 enzyme reconstitution
enzyme stability
 enzyme subunit
 methylobacterium
 molecular size
 nonhuman
 oxidation
 priority journal
 *calcium ion
 *oxidoreductase: EC, endogenous compound
 enzyme inhibitor
 guanidine hydrochloride
 methanol
 methanol dehydrogenase: EC, endogenous compound
 pyrroloquinolinequinone: EC, endogenous compound
 urea
 unclassified drug

L8 ANSWER 12 OF 14 MEDLINE on STN

AB Chloroplast carbonic anhydrase is dependent on a reducing environment to retain its catalytic activity. To investigate the properties of the three accessible cysteine residues of pea carbonic anhydrase, **mutants** were made in which Ala or Ser substituted for C165, C269, and C272. The **mutants** at position 165 were found to be **spectroscopically** similarly to the **wild-type**. They have a high catalytic activity, and are also sensitive to oxidation. In contrast, both C269 and C272 were found to be critical both for the structure and for the catalytic activity. All **mutants** with substitutions at either of these two positions had to be co-overexpressed with GroES/EL **chaperones** to give soluble enzyme in Escherichia coli. The $k(\text{cat})$ values were decreased by 2 and 3 orders of magnitude for the C272A and C269A **mutants**, respectively, and the K_m values were increased approximately 7 times. However, the binding of the inhibitor ethoxycarbonyl was only slightly weakened. The near-UV CD spectra were found to be changed in both sign and intensity compared to that of the **wild-type**, and the far-UV spectra indicate some loss of alpha-helix structure. Moreover, the quaternary structure was changed from the **wild-type** octameric to tetrameric in these **mutants**. The results indicate that mutation of either of these cysteines causes minor structural changes around at least one of the two tryptophans of the subunit. Furthermore, the data demonstrate that C269 and C272 are involved in the interaction between subunits and are necessary for a proper structure at the tetramer-tetramer interface.

ACCESSION NUMBER: 97254614 MEDLINE

DOCUMENT NUMBER: PubMed ID: 9100024

TITLE: The sulfhydryl groups of Cys 269 and Cys 272 are critical for the oligomeric state of chloroplast carbonic anhydrase

from *Pisum sativum*.
 AUTHOR: Bjorkbacka H; Johansson I M; Skarfstad E; Forsman C
 CORPORATE SOURCE: Department of Biochemistry, Umea University, Sweden.
 SOURCE: Biochemistry, (1997 Apr 8) 36 (14) 4287-94.
 Journal code: 0370623. ISSN: 0006-2960.

PUB. COUNTRY: United States
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199705
 ENTRY DATE: Entered STN: 19970523
 Last Updated on STN: 20000303
 Entered Medline: 19970515

SO Biochemistry, (1997 Apr 8) 36 (14) 4287-94.
 Journal code: 0370623. ISSN: 0006-2960.

AB . . . environment to retain its catalytic activity. To investigate the properties of the three accessible cysteine residues of pea carbonic anhydrase, **mutants** were made in which Ala or Ser substituted for C165, C269, and C272. The **mutants** at position 165 were found to be **spectroscopically** similarly to the **wild-type**. They have a high catalytic activity, and are also sensitive to oxidation. In contrast, both C269 and C272 were found to be critical both for the structure and for the catalytic activity. All **mutants** with substitutions at either of these two positions had to be co-overexpressed with GroES/EL **chaperones** to give soluble enzyme in *Escherichia coli*. The $k(\text{cat})$ values were decreased by 2 and 3 orders of magnitude for the C272A and C269A **mutants**, respectively, and the K_m values were increased approximately 7 times. However, the binding of the inhibitor ethoxzolamide was only slightly . . . weakened. The near-UV CD spectra were found to be changed in both sign and intensity compared to that of the **wild-type**, and the far-UV spectra indicate some loss of alpha-helix structure. Moreover, the quaternary structure was changed from the **wild-type** octameric to tetrameric in these **mutants**. The results indicate that mutation of either of these cysteines causes minor structural changes around at least one of the . . .

CT . . .
 genetics

Ethoxzolamide: ME, metabolism
 Ethoxzolamide: PD, pharmacology
 Gene Expression
 Kinetics
 Molecular Weight
 Mutagenesis, Site-Directed
 *Peas: EN, enzymology
 Phosphines: PD, pharmacology
 *Protein Conformation
 Protein Structure, Secondary
 Protein Structure, Tertiary
 Spectrometry, Fluorescence

L8 ANSWER 13 OF 14 MEDLINE on STN

AB One of the major protein components of the ocular lens, alpha-crystallin, is composed of alphaA and alphaB chain subunits that have structural homology to the family of mammalian small heat shock proteins. Like other small heat shock proteins, alpha-crystallin subunits associate to form large oligomeric aggregates that express **chaperone**-like activity, as defined by the ability to suppress nonspecific aggregation of proteins destabilized by treatment with a variety of denaturants including

heat, UV irradiation, and chemical modification. It has been proposed that age-related loss of sequences at the C terminus of the alphaA chain subunit may be a factor in the pathogenesis of cataract due to diminished capacity of the truncated crystallin to protect against nonspecific aggregation of lens proteins. To evaluate the functional consequences of alpha-crystallin modification, two **mutant** forms of alphaA subunits were prepared by site-directed mutagenesis. Like **wild type** (WT), aggregates of approximately 540 kDa were formed from a tryptophan-free alphaA **mutant** (W9F). When added in stoichiometric amounts, both WT and W9F subunits completely suppressed the heat-induced aggregation of aldose reductase. In contrast, subunits encoded by a truncation **mutant** in which the C-terminal 17 residues were deleted (R157STOP), despite having **spectroscopic** properties similar to WT, formed much larger aggregates with a marked reduction in **chaperone**-like activity. Similar results were observed when the **chaperone**-like activity was assessed through inhibition of gamma-crystallin aggregation induced by singlet oxygen. These results demonstrate that the structurally conservative substitution of Phe for Trp-9 has a negligible effect on the functional interaction of alphaA subunits, and that deletion of C-terminal sequences from the alphaA subunit results in substantial loss of **chaperone**-like activity, despite overall preservation of secondary structure.

ACCESSION NUMBER: 97112991 MEDLINE
DOCUMENT NUMBER: PubMed ID: 8943244
TITLE: Cloning, expression, and **chaperone**-like activity of human alphaA-crystallin.
AUTHOR: Andley U P; Mathur S; Griest T A; Petrash J M
CORPORATE SOURCE: Department of Ophthalmology and Visual Sciences, Washington University School of Medicine, St. Louis, Missouri 63110, USA.. petrash@am.seer.wustl.edu
CONTRACT NUMBER: EY05681 (NEI)
EY05856 (NEI)
P30 EY02687 (NEI)
+
SOURCE: Journal of biological chemistry, (1996 Dec 13) 271 (50) 31973-80.
Journal code: 2985121R. ISSN: 0021-9258.
PUB. COUNTRY: United States
DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
OTHER SOURCE: GENBANK-U66582; GENBANK-U66584
ENTRY MONTH: 199701
ENTRY DATE: Entered STN: 19970219
Last Updated on STN: 19980206
Entered Medline: 19970117
TI Cloning, expression, and **chaperone**-like activity of human alphaA-crystallin.
SO Journal of biological chemistry, (1996 Dec 13) 271 (50) 31973-80.
Journal code: 2985121R. ISSN: 0021-9258.
AB . . . small heat shock proteins. Like other small heat shock proteins, alpha-crystallin subunits associate to form large oligomeric aggregates that express **chaperone**-like activity, as defined by the ability to suppress nonspecific aggregation of proteins destabilized by treatment with a variety of denaturants. . . the truncated crystallin to protect against nonspecific aggregation of lens proteins. To evaluate the functional consequences of alpha-crystallin modification, two **mutant** forms of alphaA subunits were prepared by site-directed

09/863,976

mutagenesis. Like **wild type** (WT), aggregates of approximately 540 kDa were formed from a tryptophan-free alphaA **mutant** (W9F). When added in stoichiometric amounts, both WT and W9F subunits completely suppressed the heat-induced aggregation of aldose reductase. In contrast, subunits encoded by a truncation **mutant** in which the C-terminal 17 residues were deleted (R157STOP), despite having **spectroscopic** properties similar to WT, formed much larger aggregates with a marked reduction in **chaperone**-like activity. Similar results were observed when the **chaperone**-like activity was assessed through inhibition of gamma-crystallin aggregation induced by singlet oxygen. These results demonstrate that the structurally conservative substitution. . . functional interaction of alphaA subunits, and that deletion of C-terminal sequences from the alphaA subunit results in substantial loss of **chaperone**-like activity, despite overall preservation of secondary structure.

CT Check Tags: Human; Support, Non-U.S. Gov't; Support, U.S. Gov't, P.H.S.

Aldehyde Reductase: ME, metabolism

Amino Acid Sequence

Base Sequence

Chaperonins: ME, metabolism

Circular Dichroism

Cloning, Molecular

Crystallins: CH, chemistry

*Crystallins: GE, genetics

Gene Expression Regulation

Molecular Sequence Data

Protein Conformation

CN 0 (**Chaperonins**); 0 (Crystallins); EC 1.1.1.21 (Aldehyde Reductase)

L8 ANSWER 14 OF 14 MEDLINE on STN

AB Although genetic and biochemical evidence has established that GroES is required for the full function of the molecular **chaperone**, GroEL, little is known about the molecular details of their interaction. GroES enhances the cooperativity of ATP binding and hydrolysis by GroEL (refs 4, 5) and is necessary for release and folding of several GroEL substrates. Here we report that native GroES has a highly mobile and accessible polypeptide loop whose mobility and accessibility are lost upon formation of the GroES/GroEL complex. In addition, lesions present in eight independently isolated **mutant** groES alleles map in the mobile loop. Studies with synthetic peptides suggest that the loop binds in a hairpin **conformation** at a site on GroEL that is distinct from the substrate-binding site. Flexibility may be required in the mobile loops on the GroES seven-mer to allow them to bind simultaneously to sites on seven GroEL subunits, which may themselves be able to adopt different arrangements, and thus to modulate allosterically GroEL/substrate affinity.

ACCESSION NUMBER: 93309590 MEDLINE

DOCUMENT NUMBER: PubMed ID: 8100614

TITLE: Characterization of a functionally important mobile domain of GroES.

AUTHOR: Landry S J; Zeilstra-Ryalls J; Fayet O; Georgopoulos C; Gierasch L M

CORPORATE SOURCE: University of Texas Southwestern Medical Center, Dallas 75235-9041.

SOURCE: Nature, (1993 Jul 15) 364 (6434) 255-8.

Journal code: 0410462. ISSN: 0028-0836.

PUB. COUNTRY: ENGLAND: United Kingdom

DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)

DELACROIX

09/863,976

LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199308
ENTRY DATE: Entered STN: 19930813
Last Updated on STN: 19950206
Entered Medline: 19930805

SO Nature, (1993 Jul 15) 364 (6434) 255-8.

Journal code: 0410462. ISSN: 0028-0836.

AB Although genetic and biochemical evidence has established that GroES is required for the full function of the molecular **chaperone**, GroEL, little is known about the molecular details of their interaction. GroES enhances the cooperativity of ATP binding and hydrolysis. . . . whose mobility and accessibility are lost upon formation of the GroES/GroEL complex. In addition, lesions present in eight independently isolated **mutant** groES alleles map in the mobile loop. Studies with synthetic peptides suggest that the loop binds in a hairpin **conformation** at a site on GroEL that is distinct from the substrate-binding site. Flexibility may be required in the mobile loops.

CT . . .

DNA, Bacterial

Escherichia coli

GroEL Protein

GroES Protein

*Heat-Shock Proteins: CH, chemistry

Heat-Shock Proteins: GE, genetics

Heat-Shock Proteins: ME, metabolism

Magnetic Resonance Spectroscopy

Molecular Sequence Data

Mutation

Peptide Fragments: CS, chemical synthesis

Peptide Fragments: CH, chemistry

Protein Binding

Protein Conformation

=>